arXiv:1110.1997v1 [math.AG] 10 Oct 2011

A new geometric description for Igusa's modular form $(azy)_5$

A. Fiorentino

Abstract

The modular form $(azy)_5$ notably appears in one of Igusa's classic structure theorems as a generator of the ring of full modular forms in genus 2, being exhibited by means of a complicated algebraic expression. In this work a different description for this modular form is provided by resorting to a peculiar geometrical approach.

1. *Definitions and Notations*

The symplectic group $Sp(2g, \mathbb{R})$ acts biholomorphically and transitively on the *Siegel upper half-plane* $\mathfrak{S}_g$, namely the tube domain of complex symmetric $g \times g$ matrices with positive definite imaginary part; its action is defined by:

$$\gamma \cdot \tau := (a\tau + b) \cdot (c\tau + d)^{-1} \qquad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in Sp(2g, \mathbb{R}) \tag{1}$$

Furthermore, for each $k \in \mathbb{Z}$ the non-vanishing function:

$$D(\gamma, \tau)^k := \det(c\tau + d)^k \qquad \forall\, (\gamma, \tau) \in Sp(2g, \mathbb{R}) \times \mathfrak{S}_g$$

is a factor of automorphy, *i.e.* a holomorphic function on $\mathfrak{S}_g$ satisfying the cocycle condition:

$$D(\gamma\gamma', \tau)^k = D(\gamma, \gamma'\tau)^k\, D(\gamma', \tau)^k \quad \forall \gamma, \gamma' \in Sp(2g, \mathbb{R}) \tag{2}$$

Thanks to this property, an action of $Sp(2g, \mathbb{R})$ is also well defined on the space of holomorphic functions on $\mathfrak{S}_g$ for each $k \in \mathbb{Z}$ by means of the rule:

$$(\gamma|_k f)(\tau) := D(\gamma^{-1}, \tau)^{-k}\, f(\gamma^{-1}\tau) \tag{3}$$

The action of the so-called *Siegel modular group* $\Gamma_g := Sp(2g, \mathbb{Z})$ is particularly central to the theory of modular forms; a *Siegel modular form*, or simply a *modular* *form*, of weight $k \in \mathbb{Z}^+$ with respect to a subgroup Γ of finite index in Γ_g is a holomorphic function $f : \mathfrak{S}_g \to \mathbb{C}$ satisfying the property [1]:

$$\gamma^{-1}|_k f = f \qquad \forall \gamma \in \Gamma \tag{4}$$

Modular forms with respect to a subgroup Γ of finite index in Γ_g form a ring $A(\Gamma)$ which is positively graded by the weights.

More generally, whenever χ is a character of Γ, a modular form with weight $k \in \mathbb{Z}^+$ and character χ is a holomorphic function $f : \mathfrak{S}_g \to \mathbb{C}$ satisfying:

$$\gamma^{-1}|_k f = \chi(\gamma) f \qquad \forall \gamma \in \Gamma$$

As for subgroups of finite index in Γ_g, they are, in fact, characterized by containing for some $n \in \mathbb{N}$ the so-called *principal congruence subgroup of level N*:

$$\Gamma_g(n) = \{\gamma \in \Gamma_g \mid \gamma \equiv 1_{2g} \bmod n\}$$

which is of finite index itself and normal in Γ_g; remarkable families of such subgroups are:

$$\Gamma_g(n, 2n) := \{\, \gamma \in \Gamma_g(n) \mid \;\; diag(a^t b) \equiv diag(c^t d) \equiv 0 \bmod 2n \,\}$$

$$\Gamma_{g,0}(n) := \{\, \gamma \in \Gamma_g \mid \;\; c \equiv 0 \bmod n \,\}$$

An outstanding role in constructing modular forms is actually played by *Riemann Theta functions with characteristics*; for each $m = (m', m'') \in \mathbb{Z}^g \times \mathbb{Z}^g$ they are defined on $\mathfrak{S}_g \times \mathbb{C}^g$ by the series:

$$\theta_m(\tau, z) := \sum_{n \in \mathbb{Z}^g} \exp\left\{ {}^t\left(n + \frac{m'}{2}\right) \tau \left(n + \frac{m'}{2}\right) + 2\,{}^t\left(n + \frac{m'}{2}\right)\left(z + \frac{m''}{2}\right) \right\}$$

where $\exp(w)$ stands for $e^{\pi i w}$. Since $\theta_{m+2n}(\tau, z) = (-1)^{{}^t m' n''} \theta_m(\tau, z)$, these functions are parametrized merely by means of reduced *g-characteristics*, namely vector columns $\begin{bmatrix} m' \\ m'' \end{bmatrix}$ with $m', m'' \in \mathbb{Z}_2^g$.

Throughout this paper reduced g-characteristics will be simply referred to as g-characteristics, their set being henceforward conventionally denoted by the symbol $C^{(g)}$. For each $m \in C^{(g)}$ the *Theta constant* θ_m with g-characteristic m is defined by setting:

$$\theta_m(\tau) := \theta_m(\tau, 0)$$

The only non-vanishing Theta constants are plainly seen to be those related to even characteristics, namely characteristics m satisfying $(-1)^{{}^t m' m''} = 1$. More precisely, a parity function is defined on g-characteristics by:

$$e(m) := (-1)^{{}^t m' m''} \qquad \forall\, m = \begin{bmatrix} m' \\ m'' \end{bmatrix} \in C^{(g)}$$

---

[1] When $g = 1$ a modular form is also demanded to be holomorphic on the cusp ∞.

thus dividing them into *even* and *odd* respectively if $e(m) = 1$ or $e(m) = -1$; there are then $2^{g-1}(2^g + 1)$ distinct non-vanishing Theta constants for each $g \geq 1$, each being related to an even g-characteristic, while the remaining $2^{g-1}(2^g - 1)$ Theta constants that are associated to the odd g-characteristics are trivial functions. Henceforward, the symbols $C_e^{(g)}$ and $C_o^{(g)}$ will stand respectively for the set of even g-characteristics and the set of odd g-characteristics.

An action of the group Γ_g is well defined on $C^{(g)}$ by:

$$\gamma \begin{bmatrix} m' \\ m'' \end{bmatrix} := \left[\begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} m' \\ m'' \end{pmatrix} + \begin{pmatrix} diag(c^t d) \\ diag(a^t b) \end{pmatrix} \right] mod\, 2 \tag{5}$$

Γ_g acts both on $C_e^{(g)}$ and on $C_o^{(g)}$, the parity being preserved, and the action of the subgroup $\Gamma_g(2)$ is, in particular, trivial. The action of Γ_g on k-plets of even 2-characteristics is also worth being briefly highlighted; when $g = 2$, a group isomomorphism between $\Gamma_2/\Gamma_2(2)$ and the symmetric group S_6 is naturally defined:

$$\psi_P : \Gamma_2/\Gamma_2(2) \mapsto S_6 \tag{6}$$

by merely focusing on the action of $\Gamma_2/\Gamma_2(2)$ on the six elements of $C_o^{(2)}$. By pointing to the action of S_6 on non ordered k-plets of even 2-characteristics, the set of non-ordered 3-plets of even 2-characteristics, in particular, is found to decompose into two orbits (cf. [GS93]):

$$C_3^- = \{\, \{m_1, m_2, m_3\} \mid e(m_1 + m_2 + m_3) = -1 \,\}$$

$$C_3^+ = \{\, \{m_1, m_2, m_3\} \mid e(m_1 + m_2 + m_3) = 1 \,\}$$

with $card(C_3^-) = card(C_3^+) = 60$, while non-ordered 4-plets of even 2-characteristics decompose into three orbits C_4^+, C_4^* and C_4^- (cf. [GS93]), where, in particular:

$$C_4^- = \{\, \{m_1, \ldots, m_4\} \mid \{m_i, m_j, m_k\} \in C_3^- \;\; \forall \{m_i, m_j, m_k\} \subset \{m_1, \ldots, m_4\} \,\}$$

$$C_4^+ = \{\, \{m_1, \ldots, m_4\} \mid \{m_i, m_j, m_k\} \in C_3^+ \;\; \forall \{m_i, m_j, m_k\} \subset \{m_1, \ldots, m_4\} \,\}$$

with $card(C_4^-) = card(C_4^+) = 15$.

With reference to the actions described in (1) and (5), the transformation law for Riemann Theta function can be now outlined; for each $m \in C^{(g)}$ and $\gamma \in \Gamma_g$ one has (cf. [Ig66] or [Ig83]):

$$\theta_{\gamma m}(\gamma\tau\,, {}^t(c\tau + d)^{-1} z) = \kappa(\gamma)\, \chi_m(\gamma)\, \phi(\gamma, \tau, z)\, \det(c\tau + d)^{\frac{1}{2}}\, \theta_m(\tau, z)$$

$$\forall \tau \in \mathfrak{S}_g \quad \forall z \in \mathbb{C}^g$$

where:

1. κ is such a function that $\kappa(\gamma)^4 = exp\{Tr({}^tbc)\}$ whenever $\gamma \in \Gamma_g$ and, more particularly, $\kappa(\gamma)^2 = exp\left\{\frac{1}{2}Tr(a - 1_g)\right\}$ whenever $\gamma \in \Gamma_2(2)$;
2. $\chi_m(\gamma) = \exp\{2\xi_m(\gamma)\}$ with:
$$\begin{aligned}\xi_m(\gamma) = &-\frac{1}{8}({}^tm'{}^tbdm' + {}^tm''{}^tacm'' - 2{}^tm'{}^tbcm'')+\\ &-\frac{1}{4}{}^tdiag(a{}^tb)(dm' - cm'')\end{aligned}$$
3. $\phi(\gamma, \tau, z) = \exp\left\{\frac{1}{2}{}^tz\,[(c\tau + d)^{-1}c]\,z\right\}$;
4. The branch of $\det(c\tau + d)^{\frac{1}{2}}$ is the one whose sign is positive whenever $Re\tau = 0$;

Since $\phi|_{z=0} = 1$, Theta constants transform as follows:

$$\theta_{\gamma m}(\gamma\tau) = \kappa(\gamma)\,\chi_m(\gamma)\det(c\tau + d)^{\frac{1}{2}}\theta_m(\tau)$$

the product $\theta_m\theta_n$ of two Theta constants being thus found to be a modular form of weight 1 with respect to $\Gamma_g(4, 8)$.

Second order Theta constants can be also defined by setting for each $m' \in \mathbb{Z}_2^g$:

$$\Theta_{m'}(\tau) := \theta_{\left[\begin{smallmatrix} m' \\ 0 \end{smallmatrix}\right]}(2\tau, 0) = \theta_{\left[\begin{smallmatrix} m' \\ 0 \end{smallmatrix}\right]}(2\tau)$$

Whenever $\gamma \in \Gamma_{g,0}(2)$, a remarkable transformation law holds for second order Theta constants:

$$\Theta_{m'}(\gamma\tau) = \theta_{\left[\begin{smallmatrix} m' \\ 0 \end{smallmatrix}\right]}(\tilde{\gamma}2\tau) = \kappa(\tilde{\gamma})\chi_{\left[\begin{smallmatrix} m' \\ 0 \end{smallmatrix}\right]}(\tilde{\gamma})\det(c\tau + d)^{\frac{1}{2}}\Theta_{m'}(\tau)$$

where $\tilde{\gamma} := \begin{pmatrix} a & 2b \\ \frac{1}{2}c & d \end{pmatrix} \in \Gamma_g$.

By virtue of this formula the product $\Theta_{m'}\Theta_{n'}$ of two second order Theta constants is likewise found to be a modular form of weight 1 with respect to $\Gamma_g(2, 4)$.

The product of all the non-vanishing Theta constants is a modular form of weight $2^{g-2}(2^g + 1)$ with respect to Γ_g itself whenever $g \geq 3$. The case $g = 2$ is instead a special one, the character appearing in the corresponding transformation formula being not trivial; in this case a modular form of weight 5 only with respect to $\Gamma_2(2)$ is actually gained:

$$\chi_5 := \prod_{m \in C_e^{(2)}} \theta_m = \mu \cdot \det\begin{pmatrix} \Theta_{[0\,0]} & \Theta_{[0\,1]} & \Theta_{[1\,0]} & \Theta_{[1\,1]} \\ \frac{\partial\Theta_{[0\,0]}}{\partial\tau_{11}} & \frac{\partial\Theta_{[0\,1]}}{\partial\tau_{11}} & \frac{\partial\Theta_{[1\,0]}}{\partial\tau_{11}} & \frac{\partial\Theta_{[1\,1]}}{\partial\tau_{11}} \\ \frac{\partial\Theta_{[0\,0]}}{\partial\tau_{12}} & \frac{\partial\Theta_{[0\,1]}}{\partial\tau_{12}} & \frac{\partial\Theta_{[1\,0]}}{\partial\tau_{12}} & \frac{\partial\Theta_{[1\,1]}}{\partial\tau_{12}} \\ \frac{\partial\Theta_{[0\,0]}}{\partial\tau_{22}} & \frac{\partial\Theta_{[0\,1]}}{\partial\tau_{22}} & \frac{\partial\Theta_{[1\,0]}}{\partial\tau_{22}} & \frac{\partial\Theta_{[1\,1]}}{\partial\tau_{22}} \end{pmatrix} \qquad (7)$$

where $\mu \in \mathbb{C}^*$ is a suitable non-zero constant. The square:

$$\chi_{10} := \chi_5^2 = \prod_{m \in C_e^{(2)}} \theta_m^2 \tag{8}$$

is, though, a modular form of weight 10 with respect to Γ_2. As concerns the product of all the second order Theta constants:

$$P_g := \prod_{m' \in \mathbb{Z}^g} \Theta_{m'} \tag{9}$$

this one is seen to be a modular form of weight 2^{g-1} with respect to $\Gamma_g(2)$ whenever $g \geq 2$.

The ring $A(\Gamma_1)$ of modular forms with respect to Γ_1 is classically known to be generated as a $\mathbb{C}$-algebra by the Eisenstein series $E_4^{(1)}$ and $E_6^{(1)}$ respectively of weight 4 and 6. Regarding the $g = 2$ case, Igusa's structure theorem provides a set of generators (cf. [Ig64] and [Ig67]):

Theorem 1. $A(\Gamma_2) = \mathbb{C}[E_4^{(2)}, E_6^{(2)}, \chi_{10}, \chi_{12}, \chi_{35}]$
where $E_4^{(2)}$ *and* $E_6^{(2)}$ *are the Eisenstein series respectively of weight* 4 *and* 6*,* χ_{10} *is the modular form described in (8),* χ_{12} *is a modular form of weight* 12 *obtained by a suitable symmetrization:*

$$\chi_{12} = \frac{1}{2^{17}3} \sum_{\substack{\{m_{i_1},\ldots,m_{i_6}\}\ s.\ t. \\ C_e^{(2)} - \{m_{i_1},\ldots,m_{i_6}\} \in C_4^+}} \pm(\theta_{m_{i_1}} \cdots \theta_{m_{i_6}})^4$$

and $\chi_{35} := \chi_5 \cdot (azy)_5$*, where* χ_5 *is as in (7) and* $(azy)_5$ *is defined by:*

$$(azy)_5 := \frac{1}{8} \sum_{\{m_i, m_j, m_k\} \in C_3^-} \pm(\theta_{m_i}\theta_{m_j}\theta_{m_k})^{20}$$

where the signs are to be properly chosen in order to to gain the correct symmetrization.

Due to the isomorphism ψ_P described in (6), the group $\Gamma_2/\Gamma_2(2)$ admits a sole non trivial irreducible representation of degree 1, the corresponding character being as follows:

$$\chi_P(\gamma) := \begin{cases} 1 & \text{if} \quad \psi_P([\gamma]) \quad \text{is an even permutation} \\ \\ -1 & \text{if} \quad \psi_P([\gamma]) \quad \text{is an odd permutation} \end{cases} \tag{10}$$

Then, by setting $\Gamma_2^+ = Ker\chi_P$, the following structure theorem holds (cf. [Ig64]):

Theorem 2. $A(\Gamma_2^+) = \mathbb{C}[E_4^{(2)}, E_6^{(2)}, \chi_5, \chi_{12}, (azy)_5]$.

Note 1. Amid the generators of $A(\Gamma_2^+)$ described in Theorem 2, χ_5 and $(azy)_5$ are the only ones that transform with the non-trivial character χ_P under the action of the full modular group Γ_2. The function $(azy)_5$ is, in particular, the unique modular form of weight 30 with respect to Γ_2^+ admitting a non trivial character under the action of Γ_2.

The next section is devoted to provide a different expression for $(azy)_5$ by means of a remarkable geometrical construction already described in [GS93].

2. *Geometric description of the modular form* $(azy)_5$

Second order Theta constants are related to Theta constants by Riemann's addition formula. When $g = 2$, the following relations hold in particular between the ten non-trivial Theta constants and the four second order Theta constants:

$$\theta^2_{\left[\begin{smallmatrix}0&0\\0&0\end{smallmatrix}\right]} = \Theta^2_{[0\,0]} + \Theta^2_{[0\,1]} + \Theta^2_{[1\,0]} + \Theta^2_{[1\,1]} \qquad \theta^2_{\left[\begin{smallmatrix}0&0\\0&1\end{smallmatrix}\right]} = \Theta^2_{[0\,0]} - \Theta^2_{[0\,1]} + \Theta^2_{[1\,0]} - \Theta^2_{[1\,1]}$$

$$\theta^2_{\left[\begin{smallmatrix}0&0\\1&0\end{smallmatrix}\right]} = \Theta^2_{[0\,0]} + \Theta^2_{[0\,1]} - \Theta^2_{[1\,0]} - \Theta^2_{[1\,1]} \qquad \theta^2_{\left[\begin{smallmatrix}0&0\\1&1\end{smallmatrix}\right]} = \Theta^2_{[0\,0]} - \Theta^2_{[0\,1]} - \Theta^2_{[1\,0]} + \Theta^2_{[1\,1]}$$

$$\theta^2_{\left[\begin{smallmatrix}0&1\\0&0\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[0\,1]} + 2\Theta_{[1\,0]}\Theta_{[1\,1]} \qquad \theta^2_{\left[\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[1\,0]} + 2\Theta_{[0\,1]}\Theta_{[1\,1]}$$

$$\theta^2_{\left[\begin{smallmatrix}1&1\\0&0\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[1\,1]} + 2\Theta_{[0\,1]}\Theta_{[1\,0]} \qquad \theta^2_{\left[\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[0\,1]} - 2\Theta_{[1\,0]}\Theta_{[1\,1]}$$

$$\theta^2_{\left[\begin{smallmatrix}1&0\\0&1\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[1\,0]} - 2\Theta_{[0\,1]}\Theta_{[1\,1]} \qquad \theta^2_{\left[\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right]} = 2\Theta_{[0\,0]}\Theta_{[1\,1]} - 2\Theta_{[0\,1]}\Theta_{[1\,0]}$$

A quadratic form Q_m in the variables X_1, X_2, X_3, X_4 is, therefore, associated to each even 2-characteristic m:

$$m \longmapsto Q_m \qquad\qquad \text{where} \quad \theta_m^2 = Q_m(\Theta_1, \Theta_2, \Theta_3, \Theta_4)$$

Hence, a quadric V_m in the projective space $\mathbb{P}^3$ also corresponds to each even 2-characteristic m:

$$m \longmapsto V_m := V(Q_m) = \{\, [X_1, X_2, X_3, X_4] \in \mathbb{P}^3 \;\mid\; Q_m(X_1, X_2, X_3, X_4) = 0 \,\}$$

Furthermore, for each 4-plet $M \in C_4^+$ the set

$$\bigcap_{m \in M^c} V_m \subset \mathbb{P}^3$$

where M^c stands for the 6-plet of even 2-characteristics being complementary in $C_e^{(2)}$ to M, contains exactly four points (cf. [GS93]). A configuration of four hyperplanes in $\mathbb{P}^3$ is thus uniquely determined for any $M \in C_4^+$, each hyperplane being characterized by passing through all except one of the four points contained in the set $\bigcap_{m \in M^c} V_m$. Therefore, a collection of four linear forms describing these four hyperplanes is found to be associated to each $M \in C_4^+$:

$$\begin{cases} \psi_1^M := \psi_1^M(X_1, X_2, X_3, X_4) \\ \psi_2^M := \psi_2^M(X_1, X_2, X_3, X_4) \\ \psi_3^M := \psi_3^M(X_1, X_2, X_3, X_4) \\ \psi_1^M := \psi_1^M(X_1, X_2, X_3, X_4) \end{cases}$$

Hence, a tetrahedron T_M in the projective space $\mathbb{P}^3$ is uniquely determined by each 4-plet $M = (m_1, m_2, m_3, m_4) \in C_4^+$, the set $\bigcap_{m \in M^c} V_m$ being in fact the set of its vertices.

A remarkable holomorphic function can be associated to each $M \in C_4^+$ by means of the functions ψ_i^M:

$$F_M(\tau) := \tilde{F}_M(\, \Theta_{[0\,0]}(\tau), \Theta_{[0\,1]}(\tau), \Theta_{[1\,0]}(\tau), \Theta_{[1\,1]}(\tau)\,)$$

$$\text{where} \quad \tilde{F}_M := \textstyle\prod_{i=1}^4 \psi_i^M$$

In particular, the 4-plet

$$M_0 := \left\{ \begin{bmatrix} 00 \\ 00 \end{bmatrix}, \begin{bmatrix} 00 \\ 01 \end{bmatrix}, \begin{bmatrix} 00 \\ 10 \end{bmatrix}, \begin{bmatrix} 00 \\ 11 \end{bmatrix} \right\} \in C_4^+$$

is such that:

$$\bigcap_{m \in M_0^c} V_m = \{[1,0,0,0], [0,1,0,0], [0,0,1,0], [0,0,0,1]\}$$

the faces of the corresponding tetrahedron T_{M_0} being thus described by:

$$\psi_1^{M_0} = X_1; \quad \psi_2^{M_0} = X_2; \quad \psi_1^{M_0} = X_3; \quad \psi_4^{M_0} = X_4;$$

Hence:

$$F_{M_0}(\tau) = \prod_{m' \in \mathbb{Z}^2} \Theta_{m'} = P_2(\tau)$$

and F_{M_0} is, therefore, a modular form of weight 2 with respect to $\Gamma_2(2)$ (cf. (9)).

Lemma 1. *The group $\Gamma_{2,0}(2)$ is the stabilizer St_{M_0} of the 4-plet M_0.*

Proof. On the one hand $\Gamma_{2,0}(2) \subset St_{M_0}$ by definition; on the other hand, $\gamma \in St_{M_0}$ implies $diag({}^t cd) - cm'' \equiv 0 \bmod 2$ for each $m'' \in \mathbb{Z}_2^2$, hence $\gamma \in \Gamma_{2,0}(2)$. □

As C_4^+ is an orbit, Lemma 1 self-evidently implies the following:

Corollary 1. $[\Gamma_2 : \Gamma_{2,0}(2)] = 15$

The map $[\gamma] \mapsto T_{\gamma M_1}$ is thus a bijection between $\Gamma_2/\Gamma_{2,0}(2)$ and the collection $\{T_M\}_{M\in C_4^+}$ of the tetrahedrons; the product of all the images of F_{M_0} under the action of the representatives of the fifteen cosets of $\Gamma_{2,0}(2)$ in Γ_2 is then a proper candidate to be focused on. However, the behaviour of F_{M_0} under the action of $\Gamma_{2,0}(2)$ is due to be investigated foremost.

Proposition 1. *Let χ_P be the character introduced in (10); then, with reference to the action in (3), one has:*

$$\eta^{-1}|_2 F_{M_0} = \chi_P(\eta)\, F_{M_0} \qquad\qquad \forall \eta \in \Gamma_{2,0}(2)$$

Proof. One only needs to check that χ_P is the very character involved in the transformation law for F_{M_0} under the action of $\Gamma_{2,0}(2)$. Since one has:

$$F_{M_0}(\eta_0\tau) = -F_{M_0}(\tau) \qquad\qquad \text{for} \quad \eta_0 = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in \Gamma_{2,0}(2)$$

then F_{M_0} transforms with a non-trivial character χ under the action of $\Gamma_{2,0}(2)$; the character χ is of course trivial on $\Gamma_{2,0}(2)^+ := Ker\chi \supset \Gamma_2(2)$, thus extending to a non-trivial character of Γ_2 (cf. [Ib91] Appendix), which is well defined on $\Gamma_2/\Gamma_2(2)$; hence χ extends to χ_P, this one being the sole non-trivial character of $\Gamma_2/\Gamma_2(2)$. □

A holomorphic function is well defined for any fixed $\gamma \in \Gamma_2$:

$$\varphi_\gamma(\tau) := \chi_P(\gamma)^{-1}(\gamma^{-1}|_2 F_{M_0})(\tau)$$

In particular, for a fixed choice $\gamma_1, \cdots \gamma_{15}$ of the representatives of the cosets of $\Gamma_{2,0}(2)$ in Γ_2 a notable holomorphic function is defined:

$$\varphi(\tau) := \prod_{i=1}^{15} \varphi_{\gamma_i}(\tau)$$

Proposition 2. *The function φ does not depend on the choice of the coset representatives.*

Proof. If $\gamma_1, \ldots, \gamma_{15}$ and $\gamma'_1, \ldots, \gamma'_{15}$ are two different choices for the representatives of the cosets of $\Gamma_{2,0}(2)$ in Γ_2, then there exists a permutation j of the indices such that for each $i = 1, \ldots, 15$ one has $\gamma'_{j(i)} = \eta_i\gamma_i$ with $\eta_i \in \Gamma_{2,0}(2)$. Hence, by

(2) and Proposition 1:

$$\begin{aligned}\prod_{j=1}^{15}\varphi_{\gamma'_j}(\tau) = \prod_{i=1}^{15}\varphi_{\gamma'_{j(i)}}(\tau) = \prod_{i=1}^{15}\varphi_{\eta_i\gamma_i}(\tau) = \prod_{i=1}^{15}\chi_P(\eta_i\gamma_i)^{-1}((\eta_i\gamma_i)^{-1}|_2F_{M_0})(\tau) = \\ = \prod_{i=1}^{15}\chi_P(\eta_i\gamma_i)^{-1}\,D(\eta_i\gamma_i,\tau)^{-2}\,\chi_P(\eta_i)\,D(\eta_i,\gamma_i\tau)^2\,F_{M_0}(\gamma_i\tau) = \\ = \prod_{i=1}^{15}\chi_P(\gamma_i)^{-1}\,D(\gamma_i,\tau)^{-2}\,F_{M_0}(\gamma_i\tau) = \prod_{i=1}^{15}\varphi_{\gamma_i}(\tau)\end{aligned}$$

□

Theorem 3. *φ is a modular form of weight 30 with respect to Γ_2 with character χ_P.*

Proof. The transformation formula $\varphi(\gamma\tau) = \chi_P(\gamma)\,D(\gamma,\tau)^{30}\varphi(\tau)$ has to be proved whenever $\gamma \in \Gamma_2$. Let thus $\gamma_1,\dots,\gamma_{15}$ be a fixed collection of coset representatives of $\Gamma_{2,0}(2)$ in Γ_2; then, for each $\gamma \in \Gamma_2$ one has:

$$\begin{aligned}\varphi(\gamma\tau) = \prod_{i=1}^{15}\varphi_{\gamma_i}(\gamma\tau) = \prod_{i=1}^{15}\chi_P(\gamma_i)^{-1}D(\gamma_i,\gamma\tau)^{-2}\,F_{M_0}(\gamma_i\gamma\tau) = \\ = \prod_{i=1}^{15}\chi_P(\gamma_i)^{-1}\,D(\gamma_i\gamma,\tau)^{-2}\,D(\gamma,\tau)^2\,F_{M_0}(\gamma_i\gamma\tau) = \\ = D(\gamma,\tau)^{30}\prod_{i=1}^{15}\chi_P(\gamma)\,\varphi_{\gamma_i\gamma}(\tau) = \\ = \chi_P(\gamma)^{15}D(\gamma,\tau)^{30}\prod_{i=1}^{15}\varphi_{\gamma_i\gamma}(\tau) = \chi_P(\gamma)\,D(\gamma,\tau)^{30}\varphi(\tau)\end{aligned}$$

the last equality being due to Proposition 2. □

Corollary 2. *There exists $\lambda \in \mathbb{C}^*$ such that $\varphi = \lambda\,(azy)_5$.*

Proof. By Theorem 3 φ is a modular form of weight 30 with respect to Γ_2^+ transforming with a non-trivial character under the action of Γ_2; hence, the thesis follows by Theorem 2 and Note 1. □

Such an expression for the modular form $(azy)_5$ has been recently found by Gehre and Krieg in a different way by means of quaternionic Theta constants (cf. [GK10]).

References

[GS93] *B. Van Geemen, D. Van Straten*, The cusp forms of weight 3 on $\Gamma_2(2,4,8)$, Mathematics of Computation 61 (1993)

[Ib91] *T. Ibukiyama*, On Siegel Modular Varieties of Level 3, International Journal of Mathematics, Volume 2 (1991)

[Ig64] *J. Igusa*, On Siegel modular forms of genus two (II), American Journal of Mathematics 86 (1964)

[Ig66] *J. Igusa*, On the graded ring of theta-constants (II), American Journal of Mathematics 88 (1966)

[Ig67] *J. Igusa*, Modular forms and projective invariants, American Journal of Mathematics 89 (1967)

[Ig83] *J. Igusa*, Multiplicity one theorem and problems related to Jacobi's formula, American Journal of Mathematics 105 (1983)

[GK10] *D. Gehre, A. Krieg*, Quathernionic Theta constants, Arch. Math. 94 (2010)

[SM94] *R. Salvati Manni*, Modular varieties with level 2 Theta strucutre, American Journal of Mathematics 116 (1994) n.6